

## ROLE OF MICRO ORGANISMS IN TEMPLE WASTES MANAGEMENT ASSISTED BY *EUDRILLUS EUGENIA*

Karthika Arumugam<sup>1</sup>, Anbuarasi Venkatachalam<sup>1</sup>, Vasanthy Muthunarayanan <sup>\*1</sup>, Seetha Devi Ganesan<sup>1</sup>, Swabna Vivek<sup>1</sup> and Susila Sugumar<sup>1</sup>

<sup>1</sup>Department of Environmental Biotechnology, Bharathidasan University, Tiruchirappalli, Tamil Nadu, India

\*Corresponding Author Tel: +91-431-2407088 (Office), + 94430 22816 ( mobile)

E-Mail: vasanthy02@yahoo.co.in; drvasanthy02@gmail.com

**ABSTRACT** The increasing solid waste generation rates and their high collection cost were some of the major problems in India. This paper deals with the effective management of Temple wastes. The temple waste consists of coconut, leaves, flowers, fruits etc which are of biodegradable in nature. It also includes milk products, grains and water, which could facilitate the growth of microorganisms. Such biodegradable temple wastes are released as such into the water bodies (or) dumped haphazardly which causes environmental pollution. Recycling of such temple waste is performed through vermitechology employing *Eisenia fetida*. Three types of waste were prepared. They were as follows: T<sub>A</sub> (control), T<sub>B</sub> (Temple waste+ Cow dung), T<sub>C</sub> (Temple waste+ Cowdung+ Microbial inoculum) in different proportions in the 1 kg capacity Reactor vessels. The physico-chemical parameters like pH, Electrical Conductivity, Total Kjeldahl nitrogen, Total Organic Carbon, Total Organic Matter and Carbon-Nitrogen ratio and phosphorus were tested in the vermicompost. Bacterial strains isolated from vermicompost and identified and are used as an inoculum to check whether they could enhance the rate of degradation. The impact of the prepared vermicompost on chick pea seed germination was also studied. These studies showed a significant growth in terms of length and height of the plant with T<sub>C</sub> vermicompost when compared to T<sub>A</sub> and T<sub>B</sub> vermicompost. Thus, it is concluded that the vermicomposting supplemented with the bacterial cultures is comparatively efficient technology for the effective management of the temple waste in an eco friendly manner.

**Keywords:** Vermicompost, Temple waste, *Eisenia fetida*, Chick pea.

---

### INTRODUCTION

Solid waste includes items that are discarded by the public, and is now a growing problem at global, regional and local levels. In Indian cities, the average solid waste generation is nearly about 1.6 million tonnes per day, (<http://panchabuta.com/2012/10/02/india-green-it-spending-to-reach-70bn-by-2015-gartner/>). The problem is more acute in developing nations than in the developed nations (Sudhanshu Kaushik and Bishambhar Datt Joshi, 2001). So, we are in need of an useful and an effective technology for the waste management. Vermicomposting offers rapid recovery of valuable resources from biodegradable waste by producing humic like substance called vermicompost. (Pramanik et al., 2007). The waste material is decomposed by the earthworm and their gut associated microbes. These earthworms also have great capacity in assimilating heavy metals present in solid waste as pollutants (Azizi et al., 2013). Many latest studies are

available regarding the potential of earthworms for vermicomposting of solid waste (Appelhoff et al., 1998; Yan et al., 2013). During vermicomposting the soil micro flora such as bacteria, fungi, yeast and actinomycetes plays a very important role directly or indirectly. In addition, the composted materials have a clear impact on soil biological properties, such as increase in microbial biomass, soil enzymes, and soil structure (Hosseini et al., 2014; Ros et al., 2006). Several examples in the literature show that compost and vermicompost are able to enhance the growth of a wide range of plant species by the supply of nutrients (Grigatti et al., 2007). Vermicomposting not only provides nutritional elements but also shows insecticidal activity and also they were proved to contain plant growth stimulating hormones (Tomati et al., 1983). So this technology is considered as one of the best methods for managing the waste generation. Solid wastes of varied composition are seen in and around cities and also at sacred places like temples. Comparing other types of solid waste, large amount

of waste are left over by pilgrims (Gurav and pathade,2011). This causes several issues such as environmental pollution and also serves as source of inoculums for many diseases (Blackman, 1995). Astemple waste contains nutritive rich material it may act as a best raw material for the vermicomposting. In the present study, attempts were made for the conversion and utilization of temple waste in an eco friendly manner

## MATERIALS AND METHODS

### Collection of the wastes

Cow dung was collected from dairy yard at the Agriculture field near the Bharathidasan University campus, Trichy. Tamilnadu, India. Temple wastes were collected from Malaikottai temple, Trichy Tamilnadu, India and the Epigeic (or) surface dwelling earthworms *Eudrillus eugeniae* was obtained from Tamilnadu Agricultural and Research University, Madurai, Tamilnadu, India.

### Preparation of experimental media

Three type of experimental media were prepared as follows. T<sub>A</sub> contains only cow dung and temple waste which is considered as control, T<sub>B</sub> contains cow dung, temple waste and earthworms and, T<sub>C</sub> contains cow dung, temple waste, earthworms and microbial inoculum. These feed mixtures were subjected for Vermicomposting in reactor vessels with the following dimensions (diameter- 40 cm, depth -9cm). The mixtures were turned over manually for 15 days in for acclimatization process. Further, 30 non-ciliated earthworms were introduced into each reactor vessels. Also 50 ml of microbial Inoculum was added to the T<sub>C</sub> media. The moisture content was maintained at 60-80% throughout the study period by periodically sprinkling of adequate quantities of water.

### Physico chemical parameter Analysis

Physicochemical analysis was performed by collecting the sample once in a week. The physico chemical parameters like Total Kjeldahl Nitrogen is analysed (Singh and pradhan, 1987) method Total Organic Carbon (Walkely and Black 1934), Total Organic Matter and Carbon-Nitrogen ratio and pH,

Electrical Conductivity, Phosphorus were analyzed (Tandon ,2009)

### Enumeration and isolation of microorganism

The bacterial population of the vermicompost was estimated by serial dilution technique (Travors and cook, 1992). Identification of the bacterial strains was performed by conducting Biochemical tests like gram staining, indole production test, methyl red test, voges proskauer test, triple sugar iron test, catalase test, oxidase test, citrate production test, urease test.( Benson ,1990).

### Seed germination

The chick pea seeds were selected for the seed germination studies. Five seeds were selected and surface sterilized with 0.1% mercuric chloride for 1 to 2 min followed by repeated rinsing with distilled water. The percentage of germination was calculated using the following formula.

Germination % = Number of seeds germinated / total number of seeds X100

### Statistical Analysis


One-way analysis of variance (ANOVA) was done to determine any significant difference among the parameters analyzed, at 0.05% level of significance. (SAS Procedures Guide, 1990)

## RESULTS AND DISCUSSION

The physico-chemical parameters were analysed from the prepared vermicompost and it is figuratively represented in Figures 1-7.

### pH

The pH range of 6.0-8.5 is found to be suitable for the soil in order to ensure compatibility with most plants (Hogg et al., 2002). As evident from **Fig.1**, there were changes in the pH throughout the vermicomposting process. The pH was in the range between 6 to 8 in all the 3 different types. The pH shift has been reported due to mineralization of nitrogen and phosphorus into nitrites/ nitrates and orthophosphate (Ndegwa et al., 2000)




**Figure 1.** Comparison Chart Of Ph Different Types Of Vermicompost

**Electrical Conductivity**

At the initial stage, the EC values were in the range of  $12.1 \pm 0.002$ ,  $11.3 \pm 0.003$ ,  $10.12 \pm 0.003$  in 3 different types. It has increased significantly in final

vermicompost and was in the range of  $19.15 \pm 0.06$ ,  $19.1 \pm 0.035$ ,  $17.7 \pm 0.12$ . (**Fig. 2.**) The increase in EC may be due to loss of organic matter and release of different mineral salts in available forms such as phosphate, ammonium, potassium, etc (Kaviraj and Sharma, 2003)



**Figure 2.** Comparison Chart Of Electrical Conductivity Different Types Of Vermicompost

**Total organic Carbon**

Total organic carbon (TOC) in all 3 different ratios of temple waste has reduced when compared to their initial levels. (Kaushik and Garg, 2003)

reported 20-45% loss of TOC in the form of  $CO_2$  from different industrial sludge's during vermicomposting. In temple waste the TOC reduction was in range of  $21.04 \pm 0.01$  to  $16.66 \pm 0.06$  in  $T_A$ ,  $21.5 \pm 0.06$  to  $17.2 \pm 0.06$  in  $T_B$  and  $21.4 \pm 0.06$

to  $14.4 \pm 0.06$  in  $T_C$  (Fig.3). It is well known that the earth worms modify the substrate condition which consequently promotes the carbon losses from the substrate through microbial respiration in the form

of  $CO_2$  and even through mineralization of organic matter and hence leading to the reduction in the TOC content has been noticed.




Figure 3. Comparison Chart Of Toc Of Different Types Of Vermicompost

#### Total Kjeldahl Nitrogen

The Total Kjeldahl nitrogen content has increased by 57.6%, 65% and 73.9% in all the three different types of treatment (Fig 4). Losses of organic carbon

might be responsible for nitrogen addition in the form of mucus nitrogenous excretory substances, growth stimulatory hormones and enzymes from the gut of the earth worms (Viel et al., 1987)





Figure 4. Comparison Chart Of Tkn Of Different Types Of Vermicompost

**C: N ratio**

C: N ratio is one of the most widely used indicators to measure the vermicompost maturation, which is reported to decrease sharply during vermicomposting process (Suther, 2008). During the temple waste degradation process, the reduction

in C: N was in the range of 49.2% in  $T_A$ , 53.04% in  $T_B$  and 61.3% in  $T_C$  (**Fig 5**). According to (Sensei,1989 )the decline in C: N ratio <40% indicates an advance degree of organic matter stabilization and reflect a satisfactory degree of organic wastes degradation




**Figure 5.** Comparison Chart Of C:N Ratios Of Different Types Of Vermicompost

**Phosphorus**

Total phosphorus (TP) was greater in final vermicompost than in the initial feed mixture. In

this study **Fig 6** shows the increase of 34%, 35%, 37% in TP in all three types. This indicates that the degradation activity is performed by the phosphate solublizing microbes present in earthworm gut. (Satchell and Martin, 1984).




**Figure 6.** Comparison Chart Of Phosphorus Different Types Of Vermicompost

**Total Organic Matter**

Many types of organic wastes were found to be readily decomposed by soil microbes. The main product of decomposition of organic matter is carbon dioxide and water (Kaushik and Garg,

2003). Thus the decomposition of organic matter reduces the amount of TOM but leaves the compost enriched with nitrogen. The reductions in TOM were in the range of 39%, 40%, and 59% (Fig 7) with respect to all the three different types of temple waste.



**Figure 7.** Comparison Chart Of Tom Of Different Types Of Vermicompost

**Statistical analysis**

**Table 1** represent the statistical data for the physico chemical parameters. The results were statistically analyzed at 0.05 levels using one way analysis of variance (ANOVA) using SPSS version 16.0. In the

results since the p value is <0.05, the variance between the three different types is significant. Therefore it is concluded that the samples mean of T<sub>A</sub>, T<sub>B</sub> and T<sub>C</sub> are different from each other significantly.

**Table 1.** Statistical representation of Physicochemical Analysis of Vermicompost using One way ANOVA

S.No	Parameters	F-Value	P-value	Df	Interpetation
1.	TOC	2.209	0.166	2,9	Not Significant
2.	TKN	4.384	0.097	2,9	Significant
3.	C:N	1.677	0.24	2,9	Not significant
4.	TOM	2.224	0.164	2,9	Not Significant
5.	pH	1.856	0.211	2,9	Not significant
6.	EC	5.962	0.12	2,9	Significant
7.	Phosphorous	0.605	0.567	2,9	Not significant
8.	Moisture	1.856	0.211	2,9	Not significant

The significance of variance, ie., p-value, given under the head sig. is 0.000. Since the significant value is less than 0.05 (p<0.05), the variance between the mean value are not significantly different. This reflects the stabilization of the compost. But in cause of TKN and EC the varicance is significantly different.

**Isolation and identification of micro organisms**

During Vermicomposting, microorganisms are considered as the primary decomposers. These microorganisms include bacteria, fungi, actinomycetes and protozoa. They break up waste

debris and facilitate the decomposition (Haritha et al., 2009). In our study with respect to **Table 2** the microbial activity was found to be higher in the initial stages THTC (Too High To Count) due to enrichment of nutrient, and subsequently reduced during the final stages due to the degradation of the waste.

**Table 2.** Total Bacterial count of Vermicomposts

S.No	Compost	Total Heterotrophic bacterial population(CFU/ml)	
		Before	After
1.	T <sub>A</sub> (control)	THTC	84±0.5 X 10 <sup>-4</sup>
2.	T <sub>B</sub> (temple waste+Cowdung)	THTC	106±0.5 X 10 <sup>-4</sup>
3.	T <sub>C</sub> (temple waste+Cowdung + microbial inoculation)	THTC	158±0.5 X 10 <sup>-4</sup>

This will affect the complete degradation of the waste. To overcome this we identified the microbes present in the vermicompost and re inoculate it manually into the vermin bin to enhance the degradation process. **Table 3** shows the Biochemical Characterization of the isolated microorganism. The bacterial genus identified from

vermicompost is *Shigella*, *Proteus*, *Salmonella* and *Citrobacter*. The isolated organisms were grown as a consortium in 50ml nutrient broth and then the culture was inoculated into the T<sub>C</sub> reactor vessel for enhancement purpose. Plates 1 and 2 represent the waste before and after the vermicomposting of T<sub>C</sub> waste.

**Table 3.** Identification of heterotrophic bacteria

S.No	Isolated colonies	Shape	Gram staining	Motility	Indole	MR	VP	TSI	Citrate	Oxidase	Catalase	Urease	Name of the genus
1.	S1	Rods	Gram negative	Non Motile	-ve	+ve	-ve	-ve	-ve	+ve	-ve	-ve	<i>Shigella</i>
2.	S2	Rods	Gram negative	Motile	-ve	+ve	-ve	+ve	+ve	-ve	+ve	+ve	<i>Proteus</i>
3.	S3	Cocci	Gram positive	Non Motile	-ve	+ve	-ve	+ve	-ve	-ve	+ve	+ve	<i>Micrococcus</i>
4.	S4	Rods	Gram negative	Motile	-ve	+ve	-ve	-ve	-ve	-ve	+ve	-ve	<i>Salmonella</i>
5.	S5	Cocci	Gram positive	Non Motile	-ve	+ve	-ve	+ve	-ve	-ve	+ve	+ve	<i>Micrococcus</i>
6.	S6	Rods	Gram negative	Motile	-ve	+ve	-ve	+ve	+ve	+ve	+ve	+ve	<i>Citrobacter</i>

Initial  
Temple waste




Final  
Vermicompost

Plates 1 & 2 -Preparation of vermicompost during experiment

**Seed germination**

Seedling stage is the most sensitive stage in the life cycle of plant and hence it is susceptible to physical and chemical adversities (Copper et al., 1998). In the present germination studies, chick pea seeds were treated for 7 days with the three different types of vermicompost. The morphological growth parameters such as germination percentage and


seedling length were taken into consideration. Fig 10 represent the effect of vermicompost on seed germination. With reference to Fig 9, the highest germination percentage was observed in T<sub>C</sub> (92%) compared to T<sub>A</sub> (58%). which may be due to activities of microbes. Also the growth and length pattern of Shoot got increased in the plate where vermicompost is used.



**Figure 8.** Effect of application of different types of vermicompost in shoot length of chick pea

As evident from Fig 8 the hieghtest Shoot length was recorded in the T<sub>C</sub> (7.8cm). The marked difference in the shoot length of chick pea may be due to the presence of growth promoting hormones which could be present in the vermicompost prepared by using temple waste along with cow dung and microbial inoculum. Similarly, Arancon et al., (2004) found that maximum benefit from vermicompost is obtained when it constitutes between 10 to 40% of the growing

medium Neilson, (1965) and Tomati, (1988) have also reported that the vermicompost contains growth promoting hormone ‘auxins’, ‘cytokinins’ and flowering hormone ‘gibberellins’ secreted by earthworms. Similarly good growth of three flowering plants such as *Mirabilis jalapa*, *Calendula officinalis*, *Clitoria ternatea* were reported by Bhat and Limaye, (2012) when vermicompost was applied to the soil.



**Figure 9.** Effect of application of different types of vermicompost in percentage of seed germination of chick pea


### CONCLUSION

In this study, the temple waste along with cow dung and microbial inoculums works as an excellent raw material for vermicomposting using *Eudrillus eugeniae*. This study concludes that the degradation was enhanced only after the addition of the microbial inoculum. This work proves that the major role of degradation was played by the microorganisms present in the vermicompost. Also the vermicompost prepared from temple waste act as a good fertilizer for the plantation, which were confirmed by seed germination studies. Hence this ecofriendly method of temple waste management

could be extended for the temples to minimize the solid waste pollution. This method greatly help us to potentially convert the temple waste into value added product in a shorter time effectively.

### ACKNOWLEDGEMENT

The author's acknowledge financial support provided by **Ministry of Environment and Forests, New Delhi, India** towards the sanction of Major Research Project. We thank **UGC-NON SAP, New Delhi and DST-FIST, New Delhi, India** for providing the instrumental facilities for the successful completion of the study.



**Figure 10.** Effect of different types of vermicompost on seed germination and seedling length of chick pea  
Plate 1-Water; Plate 2-Control; Plate 3: Temple Waste; Plate 4: Temple waste +Inoculum

## REFERENCES

1. Appelholf, M., Edward, C.A., Neuhauser, E.F., 1998. Domestic vermicomposting systems Earthworm waste. *Environ Mange*, 157-161.
2. Arancon, N.Q., Edwards, C.A., Bierman, P., Welch, C., Metzger J.D., 2004. Influences of vermicomposts on field strawberries: Effects on growth and yields. *Bioresource Techn*, 93,145-153.
3. Azizi, A.B., Lim, M.P.M., Noor, Z.M., Abdullah, N., 2013. Vermiremoval of heavy metal in sewage sludge by utilising *Lumbricus rubellus*. *Ecotoxicology and Environmental Safety*, 90, 13–20.
4. Benson, H.J., 1990. Microbiological applications-a laboratory manual in general microbiology. *WMC Brown Publishers*.
5. Bhat ,M.R., Limaye, S.R., 2012. Nutrient status and plant growth promoting potential of prepared vermicompost. *International journal of Environmental Sciences*,3(1), 312 -321.
6. Blackman ,C.W., 1995. Basic Hazardous waste management. *Lewis Publishers*, Newyork, USA.
7. Cooper, R., Chunhua, L., Fisher, D., 1998. Influence of humic substances on rooting and nutrient content of creeping bentgrass. *Crop Sci*, 38, 1639-1644.
8. Grigatti, M., Giorgonni, M.E., Ciavatta, C., 2007. Compost-based growing media: influence on growth and nutrient use of bedding plants. *Bioresource Technology*, 98, 3526-3534
9. Gurav,M.V., Pathade,G.R., 2011. Production of Vermicompost from Temple Waste (Nirmalya): A Case Study .*Universal Journal of Environmental Research and Technology*, 1(2), 182-192.
10. Haritha Devi, S., Vijayalakshmi, K., Pavana Jyotsna, K., Shaheen, S.K., Jothi, K., Surekha rani, M., 2009. Comparative assessment in enzyme activities and microbial populations during normal and vermicomposting. *Journal of Environmental Biology*, 30(6), 1013-1027.
11. Hogg, D., Barth, J., Favoino, E., Centemero, M., Caimi, V., Amlinger, F., Devliegher, W., Brinton, W., Antler, S., 2002. Review of Compost Standards in Australia. *Banbury, Reino Unido*. 11.
12. Hossein, M., Mohammad, F., Alireza, S., Mohammad, A., Mohsen, N., Shila, A., Khashayar, R., 2014. Effect of vermicompost on plant growth and its relationship with soil properties. *International Journal of Farming and Allied Sciences*, 3-3, 333-338.
13. Kaushik, P.,Garg, V.K.,2003.ermicomposting of mixed solid textile mill sludge and cow dung with the epigeic earthworm *Eisenia fetida*. *Bioresour Technol*, 90, 311-316.
14. Kaviraj., Sharma, S., 2003. Municipal solid waste management through vermicomposting employing exotic and local species of earthworms. *Bioresour. Technol*, 90, 169-173.
15. Ndegwa, P.M., Das, K.C., Thompson, S.A., 2000. Effects of stocking density and feeding rat on vermicomposting of biosolids. *Bioresource Technology*, 71(1), 5-12.
16. Neilson, R.L., 1965. Presence of Plant Growth Substances in Earthworms, Demonstrated by the Paper Chromatography and Went Pea Test, *Nature (Lond)*, 208, 1113-1114.
17. Pramanik, P., Ghosh, G.K., Chosal, P.K., Banik,P., 2007. Changes in organic - C, N, P and K and enzyme activities in vermicompost of biodegradable organic wastes under liming and microbial inoculants. *Biores. Technol*, 98, 2485-2494
18. Ros, M., Pascual, J.A., García, C., Hernández, M.T., Insam, H., 2006. Hydrolase activities, microbial biomass and bacterial community in a soil after long-term amendment with different composts. *Soil Biology and Biochemistry*, 38, 3443-3452

19. SAS Institute, 1990. SAS Procedures Guide, Version 6, third ed. SAS Institute, Cary.
20. Satchell, J.E., Martin, K., 1984. Phosphatase activity in earthworm faeces. *Soil Biol. Biochem*, 2, 191-194.
21. Senesi, N., 1989. Composted material as organic fertilizer. *The sci. Total Environ*, 81(82), 521-524.
22. Singh, R., Pradhan, K., 1981. Determination of nitrogen and protein by Kjeldahl method Forage *Evaluation Science*, Pvt : Publishers 23.
23. Sudhanshu, K., Bishambhar, D.J., 2001. A comparative study of solid waste generation at Mansa Devi and Chandi Devi temples in Shivalik foothills, during the Kumbh mela 2010. Department of Zoology and Environmental Science, Faculty of life science, *Gurukula Kangri Vishwavidyalaya in Haridwar*, Uttarakhand, India.
24. Suthar, S., 2008. Bioconversion of post harvest crop residues and cattle shed manure into value added products using earthworm *Eudrilus eugeniae*. *Ecol. Eng*, 32, 206-214.
25. Tandon, H.L.S., 2009. Methods of Analysis of Soils Plants, Waters, Fertilisers and Organic Manures. Fertiliser Development and consultation Organisation, New Delhi. India.
26. Tomati, V., Grappelli, A., Galli, E., 1988. The Hormone like Effect of Earthworm Casts on Plant Growth. *Biology and Fertility of Soils*, 5, 288-294.
27. Tomati, V., Grapelli, A., Galli, E., Rossi, W., 1983. Fertilizers from vermiculture as an option for organic waste recovery. *Agrochemica*, 2, 244-251.
28. Travors, J.T., Cook, S., 1992. A comparison of plating media and diluents for enumeration of aerobic bacteria in a loam soil. *Journal of Microbe Methods*, 14, 271-275.
29. Viel, M., Sayag, D., Andre, L., 1987. Optimization of agricultural, industrial waste management through in-vessel composting. In: De Bertoldi M (ed) *Compost: production, quality and use*, Elsevier, Essex, 230-237.
30. Walkley, A, Black, I.A., 1934. Chromic acid titration for determination of soil organic matter. *J Soil Sci*, 63, 251.
31. Yan, Y.W., Nor A., Shamsuddin, Z.H., Muskhazli, M., Suraini, A.A., Teng, S.K. 2013 Comparison of plant nutrient contents in vermicompost from selected plant residues *African Journal of Biotechnology*, 12(17), 2207-2214.