

# Genotypic Detection of *rpoB* and *katG* Gene Mutations Associated with Rifampicin and Isoniazid Resistance in *Mycobacterium Tuberculosis* Isolates: A Local Scenario (Kelantan)

Nurul-Ain ISMAIL<sup>1</sup>, Mohd Fazli ISMAIL<sup>2</sup>, Siti Suraiya MD Noor<sup>2</sup>,  
Siti Nazrina CAMALXAMAN<sup>1</sup>

Submitted: 25 Jan 2015  
Accepted: 14 Oct 2015

<sup>1</sup> Department of Medical Laboratory Technology, Faculty of Health Sciences, Universiti Teknologi MARA, 42300, Bandar Puncak Alam, Selangor, Malaysia

<sup>2</sup> Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia

## Abstract

**Background:** Drug resistant tuberculosis (DR-TB) remains a public health issue that is of major concern on a global scale. The characterisation of clinical isolates may provide key information regarding the underlying mechanisms of drug resistance, and helps to augment therapeutic options. This study aims to evaluate the frequency of gene mutations associated with Rifampicin (RIF) and Isoniazid (INH) resistance among nine clinical isolates.

**Methods:** A total of nine drug resistant *Mycobacterium tuberculosis* clinical isolates were screened for genetic mutations in *rpoB* and *katG* using polymerase chain reaction (PCR) amplification and DNA sequencing. Genotypic analysis was performed to detect the mutations in the sequence of the target genes.

**Results:** Our findings reveal that 80% of the isolates possess mutations at codon 119 (His119Tyr) and 135 (Arg135Trp and Ser135Leu) within the *rpoB* gene; and 70% possess mutations in the *katG* gene at codon 238 with amino acid change (Leu238Arg).

**Conclusion:** Findings from this study provide an overview of the current situation of RIF and INH resistance in a hospital Universiti Sains Malaysia (HUSM) located in Kelantan, Malaysia, which could facilitate molecular-based detection methods of drug-resistant strains. Further information regarding the molecular mechanisms involved in resistance in RR-/MDR-TB should be addressed in the near future.

**Keywords:** *M. tuberculosis*, isoniazid, rifampicin, *rpoB*, *katG*, mutation

## Introduction

Tuberculosis (TB) is among the major public health concerns, and is the leading cause of morbidity and mortality related with infectious diseases worldwide. The aetiological agent is *Mycobacterium tuberculosis*, which is transmitted by airborne particles. According to World Health Organization (WHO) reports, there have been a total of 9 million TB incidents worldwide, with 480 000 cases of multi-drug resistant tuberculosis (MDR-TB) in 2013 alone (1). Needless to say, the economic impact of TB is substantial, posing immense challenges to society.

The treatment of active TB requires the administration of Isoniazid (INH) in combination with one or more first line drugs such as Rifampicin (RIF), Ethambutol (EMB), Pyrazinamide (PZA) or

Streptomycin (STR). Drug resistance and MDR-TB strains of *Mycobacterium tuberculosis* have been recognized in many parts of the world (2,3,4), which hinders the success of TB control programs. MDR-TB is defined as the resistance to at least two main first line TB drugs: RIF and INH (1). Drug resistance is generally an inevitable outcome of the inadequate, incomplete or prolonged usage of antibiotics (5). Genetic analysis of drug-resistant strains revealed that airborne transmission of undetected and untreated strains have played a key role in several outbreaks (6). In Malaysia, 277 cases of Rifampicin Resistance-/MDR-TB (RR-/MDR-TB) have been reported in 2013, compared to 141 cases in 2011, following drug sensitivity tests by the National Public Health Laboratory (NPHL) in Sg. Buloh (7). Although the number of cases is relatively low compared to other Asian

countries, the prevalence of RR-/MDR-TB is increasing. This is attributable to the increase of immigrants, resulting in endemicity (5).

Resistance is often associated with mutations in target-encoding, or related, genes. Mutations in the hypervariable region of the RNA polymerase beta-subunit encoding gene (*rpoB*), which encodes the  $\beta$  subunit of the DNA-dependent RNA polymerase (RNAP), are likely to impact its functions (8, 9), as observed in 95% of RIF resistant strains. In addition, RIF resistance may be regarded as a surrogate marker for MDR-TB, since 83% of the strains in the UK are also resistant to INH (10). The detection of a variety of mutations that involve one or several genes in INH resistant strains is highly complicated. About 60–70% of INH resistance is associated with mutations in the catalase-peroxidase encoding (*katG*) gene (11,12,13), which is the only enzyme in TB that is capable of activating INH from pro-drug to active form (8). This study aims to determine the frequencies of mutations among drug resistant strains in TB cases at Hospital Universiti Sains Malaysia (HUSM), Kelantan.

## Materials and Methods

### *Clinical isolates and drug sensitivity testing*

HUSM provided a total of nine clinical isolates of *M. tuberculosis*, as well as the H37Rv strain as the positive control. All isolates were cultured on Lowenstein-Jensen slants, and then subjected to drug sensitivity testing, which were classified as MDR-TB by the NPHL in Sg. Buloh, Malaysia. The data has been obtained from NPHL for molecular testing.

### *DNA extraction and PCR amplification*

Genomic DNA was isolated using the boiling method according to standard procedures. Its concentration (ng/ $\mu$ L) was estimated spectrophotometrically at 260 nm. DNA fragments of the *rpoB* and *katG* genes were amplified using the primer sets described by Zenteno-Cuevas et al. (3). PCR reactions were conducted in 25  $\mu$ L reaction volumes containing 300 ng/ $\mu$ L genomic DNA. The PCR cocktail contained 1 $\times$  PCR buffer, 200 $\mu$ M dNTPs (each), 1.25 units Top Taq DNA polymerase, 1.5 mM MgCl<sub>2</sub>, and 0.2  $\mu$ M of *rpoB* and *katG* primers. Each PCR consisted of 35 cycles of denaturation at 94 °C for 3 minutes; annealing at 60 °C for 30 seconds; with an extension at 72 °C for 1 minute; and a final extension step at 72 °C for 10 minutes. PCR products were subjected to electrophoretic separation using 1.5% (w/v) agarose gels containing SYBR® Safe DNA stain,

visualized under UV light.

### *DNA sequencing*

PCR products were purified using the GeneJET PCR Purification kit (Thermo Scientific, MA, USA). The H37Rv strain and a total of nine RIF and INH resistant samples were sequenced directly using the Big Dye® Terminator v.3.1 Cycle Sequencing Kit and an automated 3130xl Genetic Analyzer (Applied Biosystems, Tokyo, Japan). The obtained sequences were directly compared to previously published *rpoB* and *katG* gene sequences using the MEGABLAST search tool in the National Center for Biotechnology Information (NCBI) database. Sequence files were imported into Chromas Lite, and then assembled using DNA Baser Sequence Assembly v.3.5.4.

### *Mutational analysis*

Genotypic analysis to detect mutations responsible for resistance was performed using the Mutation Surveyor® DNA Variant Analysis software version 4.0.9 (SoftGenetics, State College, PA).

## Results

### *PCR and sequencing*

Genotypic analysis was performed for RIF and INH resistant specimens to detect mutations responsible for drug resistance by PCR amplification of the *rpoB* gene (280bp) and *katG* gene (580bp) respectively. Nine isolates (USM 01, 31, 43, 53, 54, 60, 98, 100 and 117) with known resistance patterns to first-line drugs from NPHL were then selected for sequencing, which revealed 99% similarities to reference strains in the NCBI database.

### *Mutation analysis*

Mutations within the *rpoB* and *katG* genes were analyzed by comparing the sequence of clinical isolates to the H37Rv strain, the result of which can be observed in table 1. The USM 98 and USM 100 isolates with RIF resistance revealed mutations at codon 119 (His119Tyr); USM 31 and USM 43 at codon 135 (Arg135Trp and Ser135Leu); and USM 01 at codon 90 (Asp90Val); accounting for 80% of mutations in the *rpoB* gene. No mutation was detected in two among the five RIF-resistant isolates. In contrast, 70% of mutations in the *katG* gene of isolates with INH resistance showed missense mutation at codon 238 (Leu238Arg). USM 117 revealed changes at codon 61 (Val61Gly); USM 01 at codon 247 (Gln247His); and USM 53 at codon 62 (Ala62Thr).

The nucleotide and amino acid changes observed in the RIF/INH drug-resistant isolates are shown in figure 1.

### Discussion

Genotypic methods are often deemed superior to those of traditional phenotypic in terms of both sensitivity and specificity. Given the slow growth rate and fastidious nature of *Mycobacterium tuberculosis*, identification


and drug susceptibility testing of TB is time consuming. TB diagnosis using standard culture techniques, even in the presence of automated fluid culture methods, takes an average of 14 days to complete (14). This presents a major hindrance for any TB control program, and complicates therapeutic options. The use of molecular based methods, based on the detection of mutations in specific genes associated with resistance, is more efficient and effective. Its utilization in clinical microbiology laboratories could minimize the

**Table 1:** Characteristics of nine RIF/ INH-resistant clinical isolates of *M. tuberculosis* isolates

Isolate	<i>rpoB</i> gene			<i>katG</i> gene			Drug resistance profile
	Codon	Amino acid change	Nucleotide substitution	Codon	Amino acid change	Nucleotide substitution	
USM 01	90	D → V	GAC → GTC	238	L → R	CTC → CGC	RIF
				247	Q → H	CAG → CAC	INH
USM 31	135	R → W	CGG → TGG	-	-	-	RIF
USM 43	135	S → L	TCG → TTG	238	L → R	CTC → CGC	RIF
USM 53	-	-	-	62	A → T	GCG → ACG	INH
				238	L → R	CTC → CGC	
USM 54	-	-	-	-	-	-	INH
USM 60	-	-	-	238	L → R	CTC → CGC	RIF
USM 98	119	H → Y	CAC → TAC	238	L → R	CTC → CGC	RIF
USM 100	119	H → Y	CAC → TAC	238	L → R	CTC → CGC	RIF
USM 117	-	-	-	61	V → G	GTG → GGA	INH
				238	S → A	TCG → GCG	

[-] No mutation detected.

Abbreviations: USM = Universiti Sains Malaysia; RIF = Rifampicin; INF = Isoniazid.



**Figure 1:** Chromatograms of nucleotide substitutions detected in TB positive samples. Star (\*) represents the mutation site revealed from the samples. (A) *rpoB* mutations detected in RIF resistance cases. (B) Mutation found in *katG* gene INH resistance case.

turnaround time required to diagnose cases from weeks to hours, hence, its use in this study.

Point mutations in several key genes of *Mycobacterium tuberculosis* is believed to be the major basis governing resistance. The catalase-peroxidase enzyme of TB is encoded by the *katG* gene, which is responsible for converting isoniazid to its active form, as well as for the detoxifying activity against reactive oxidative stress (15). The mutation of this gene causes loss of, or reduced, activity of the catalase-peroxidase activity required to maintain the level of oxidative protection against host antibacterial radicals (16). Together with INH, RIF is another vanguard antimicrobial drug that is commonly used in the therapy of TB since the 1980s (17). Mutations in the *rpoB* gene generally confer resistance to RIF, commonly through amino acid changes involving a compact side chain with one that is larger (e.g., Ser Leu), thus, preventing RIF access to RNAP (9). Mutations in the *rpoB* gene have a negative impact on key physiological functions often associated with treatment failure and fatal clinical outcomes (18). Both genes were analyzed concurrently using clinical isolates from local settings.

There has been little information on the molecular characterization of drug resistant TB clinical isolates in Malaysia, which is alarming, given its endemicity. In general, our findings support prior hypotheses that link the *rpoB* and *katG* genes to RIF and INH drug resistance. All of the RIF and INH mutations were characterized as single nucleotide changes. Nevertheless, the site of mutation and type of changes differ from findings reported in prior work. The most frequent mutations in the *rpoB* gene of RIF resistant strains were at codons 135 and 119; this has been observed in 80% of the isolates. We did not observe mutation 531 (Ser531Leu), which is a common documented site of *rpoB* mutation (19,14). Furthermore, while most studies report mutation at Ser315Thr (20,19,14), our findings detected changes at codon 238 (Leu238Arg), accountable for 70% of the *katG* mutation. However, this change did not contribute to INH phenotypic resistance, since it is possibly considered the most common polymorphism or low frequency mutation in the *katG* gene. We also observed changes at codon 247 (Gln247His), codon 61 (Val61Gly), and codon 62 (Ala62Thr). Two isolates (USM 31 and USM 54) did not exhibit any mutation, despite being phenotypically resistant.

Our findings are unique compared to other reports in other parts of the world, reflecting the difference of gene mutations in different

geographical locations. Discrepancies observed could be attributed to the limited number of samples, presence of other mutations which have not been documented to date, or its presence outside selected target regions in the *rpoB* and *katG* genes. For instance, INH resistance involves alterations of multiple genes other than the *katG* gene, such as *inhA*, *kasA* and *ahpC* (13), which were beyond the scope of this work. In addition, newly undefined genes may also be implicated, and many underlying mechanisms remain unexplored. Not all mutations are known, some which are of low frequency; many vary according to geographical regions, all of which convolute the task of detection based on single genes. Although molecular-based methods offer many advantages, these limitations should be taken in consideration when designing a diagnostic assay for resistant strains.

## Conclusion

The detection of mutations in resistance determining regions plays a crucial role for the rapid detection of anti-TB resistance, and could aid strategies to further explore the mechanisms of resistance. The high percentage of mutations in the *rpoB* and *katG* genes highlights their importance, and should be further investigated in order to gain a more comprehensive of the geographical distribution of resistant strains, as well as of the mechanisms underlying them.

## Acknowledgement

The authors would like to thank the Dean of the Faculty of Health Sciences, UiTM for supporting this project financially and the Department of Medical Microbiology and Parasitology Universiti Sains Malaysia, Kubang Kerian, Kelantan for providing the clinical samples and research facilities.

## Conflict of Interest

None.

## Funds

None.

## Authors' Contributions

Conception and design: NAI, SSNM, SNC  
Analysis and interpretation of the data: NAI  
Drafting of the article: NAI, SNC

Critical revision of the article for important intellectual content, statistical expertise: SNC  
Final approval of the article: SSNM, SNC  
Provision of study materials or patients: SSNM  
Administrative, technical, or logistic support: NAI, MFI, SSNM  
Collection and assembly of data: MFI

## Correspondence

Siti Nazrina Camalxaman  
BSc, MSc Biomedical Science, PhD (Virology)  
Department of Medical Laboratory Technology,  
Faculty of Health Sciences  
Universiti Teknologi MARA,  
42300, Bandar Puncak Alam,  
Selangor, Malaysia.  
Fax: +603-32584599  
Tel: + 603-32584428  
Email: sitinazrina@salam.uitm.edu.my

## References

1. World Health Organization (WHO). *Global Tuberculosis Report 2014*. Geneva (CH): WHO Press; 2014. p. 118.
2. Temple B, Ayakaka I, Ogwang S, Nabanjja H, Kayes S, Nakubulwa S, et al. Rate and Amplification of Drug Resistance among Previously-Treated Patients with Tuberculosis in Kampala, Uganda. *Clin Infect Dis*. 2008;**47**(9):1126–1134. doi: 10.1086/592252.
3. Zenteno-Cuevas R, Zenteno JC, Cuellar A, Cuevas B, Sampieri CL, Riviera JE, et al. Mutations in rpoB and katG genes in Mycobacterium isolates from the Southeast of Mexico. *Mem Inst Oswaldo Cruz*. 2009;**104**(3):46. doi: <http://dx.doi.org/10.1590/S0074-02762009000300012>.
4. Glynn J R, Whiteley J, Bifani PJ, Kremer K, van Soolingen D. Worldwide Occurrence of Beijing/W Strains of Mycobacterium tuberculosis: A Systematic Review. *Emerg Infect Dis*. 2002;**8**(8):843–849. doi: 10.3201/eid0808.020002.
5. Aziah AM. Tuberculosis in Malaysia: Combating the old Nemesis. *Med J Malaysia*. 2004;**59**(1):1–3.
6. Keshavjee S, Farmer PE. Tuberculosis, Drug Resistance and the History of Modern Medicine. *N Engl J Med*. 2012;**367**(10):931–936. doi: 10.1056/NEJMr1205429.
7. World Health Organization (WHO). *Database: Tuberculosis country profiles*. Geneva 27, Switzerland: WHO Press; 2014 [Modified 2015 Jan; cited 2015 Jan 20]. Available from: <http://www.who.int/tb/country/data/profiles/en/>.
8. Abdelaal A, El-Ghaffar HA, Zaghoul MHE, Mashad NE, Badran E, Fathy A. Genotypic detection of rifampicin and isoniazid resistant Mycobacterium tuberculosis strains by DNA sequencing: A randomized trial. *Ann Clin Microbiol Antimicrob*. 2009;**8**:4. doi: 10.1186/1476-0711-8-4.
9. Koch A, Mizrahi V, Warner DF. The impact of drug resistance on Mycobacterium tuberculosis physiology: what can we learn from rifampicin? *Emerg Microbes Infect*. 2014;**3**: e17. doi: 10.1038/emi.2014.17.
10. O’Riordan P, Schwab U, Logan S, Cooke G, Wilkinson RJ, Davidson RN, et al. Rapid Molecular Detection of Rifampicin Resistance Facilitates Early Diagnosis and Treatment of Multi-Drug Resistant Tuberculosis: Case Control Study. *PLoS ONE*. 2008;**9**(3):1–7. doi: 10.1371/journal.pone.0003173.
11. Iqbal S, Shabir I, Iqbal R, Riazudin S. Role of katG Gene in Resistant Build up Against Isoniazid (INH)-A First Line Anti-TB Drug. *Int J Agric Biol*. 2004;**6**(3): 581–584.
12. Aragon LM, Navarro F, Heiser V, Garrigo M, Espanol M, Coll P. Rapid detection of specific gene mutations associated with isoniazid or rifampicin resistance in Mycobacteria tuberculosis clinical isolates using non-fluorescent low-density DNA microarray. *J Antimicrob Chemother*. 2006;**57**:825–831. doi: 10.1093/jac/dkl058.
13. Yang S, Zhong M, Zhang Y, Wang Y. Rapid detection of rpoB and katG genes from the sputum of multidrug-resistant Mycobacterium tuberculosis by polymerase chain reaction (PCR)-direct sequencing analysis. *Afr J Microbiol Res*. 2011;**5**(26):4519–4523. doi: 10.5897/AJMR11.358.
14. Tessema B, Beer J, Emmrich F, Sack U, Rodloff A. Analysis of gene mutations associated with isoniazid, rifampicin and ethambutol resistance among Mycobacterium tuberculosis isolates from Ethiopia. *BMC Infect Dis*. 2012; **12**:37. doi:10.1186/1471-2334-12-37.
15. Valvatne H, Syre H, Kross M, Stavrum R, Ti T, Phyu S, et al. Isoniazid and rifampicin resistance-associated mutations in Mycobacterium tuberculosis isolates from Yangon, Myanmar: implications for rapid molecular testing. *J Antimicrob Chemother*. 2009;**64**(4):694–701. doi: 10.1093/jac/dkp292.
16. Somoskovi A, Parsons LM, Salfinger M. The molecular basis of resistance to isoniazid, rifampicin and pyrazinamide in Mycobacterium tuberculosis. *Respir Res*. 2001;**2**:164–168. doi: 10.1186/rr54.
17. Mitchison DA. Role of individual drugs in the chemotherapy of tuberculosis. *Int J Tuberc Lung Dis*. 2000;**4**(9):796–806.
18. Williamson DA, Roberts SA, Bower JE, Vaughan R, Newton S, Lowe O, et al. Clinical failures associated with rpoB mutations in phenotypically occult multidrug-resistant Mycobacterium tuberculosis. *Int J Tuberc Lung Dis*. 2011;**16**(2):216–220. doi: 10.5588/ijtld.11.0178.
19. Wang YC, Zhu RY, Xu YY, Zhao MQ, Liu YH, Chen JD. Molecular characterization of drug resistant Mycobacterium tuberculosis isolates in Guangdong, China. *Jpn J Infect Dis*. 2009;**62**(4):270–274.
20. Guo H, Seet Q, Denkin S, Parsons L, Zhang Y. Molecular characterization of isoniazid-resistant clinical isolates of Mycobacterium tuberculosis from the USA. *J Med Microbiol*. 2006;**55**(11):1527–1531. doi: 10.1099/jmm.0.46718-0.